

New species of sygobiotic *Belgrandiella* from north Bulgaria (Gastropoda: Hydrobiidae)

Dilian Georgiev

Department of Ecology and Environmental Conservation, University of Plovdiv, Tzar Assen Str. 24, BG-4000 Plovdiv, Bulgaria

* **E-mail:** diliangeorgiev@abv.bg

Abstract:

Georgiev, D.: *New species of sygobiotic Belgrandiella from north Bulgaria (Gastropoda: Hydrobiidae). Biologica Nyssana, 3 (2), December 2012: 105-107.*

A new stygobiotic species of *Belgrandiella* from a cave in north Bulgaria was described. Nine shells were collected by the author on 09.03.2013 at the entrance of the Water cave (Vodnata) near village of Musina, north Bulgaria, N43 13 14.9 E25 25 39.6, 198 m alt. The systematic position of the genus *Belgrandiella* was also discussed.

Key words: aquatic, Bulgaria, endemic, subterranean, snail

Introduction

The genus *Belgrandiella* Wagner, 1927 (Gastropoda: Risooidea) consists of minute snails inhabiting the springs and the subterranean waters of south-west and central Europe to the Caucasus (Kabat, Hersler, 1993; Arconada, Ramos, 2003). This is one of the most diverse genera from the Risooidea superfamily in Bulgaria having 11 species known till now (Georgiev, 2011a, Georgiev, 2013). There are two ecological groups of species from the genus: stygobiotic (*Belgrandiella hessei* Wagner, 1927, *Belgrandiella pusilla* Angelov, 1959, *Belgrandiella bulgarica* Angelov, 1972, *Belgrandiella bureschi* Angelov, 1976, *Belgrandiella stanimirae* Georgiev, 2011, *Belgrandiella maarensis* Georgiev, 2013) and spring-living species (*Belgrandiella angelovi* Pintér, 1968, *Belgrandiella zagoraensis* Glöer & Georgiev, 2009, *Belgrandiella dobrostanica* Glöer & Georgiev, 2009, *Belgrandiella bachkovoensis* Glöer & Georgiev, 2009, *Belgrandiella pandurskii* Georgiev, 2011, the last one also found in cave waters) (Wagner, 1927; Angelov, 1959, 1972, 1976; Pintér, 1968; Hubenov, 2005, 2007;

Glöer, Georgiev, 2009; Georgiev, 2011b, c, d).

In this paper I describe a new stygobiotic species of *Belgrandiella* from a cave in north Bulgaria showing that the subterranean waters of the country are still poorly known considering their fauna and many new species could be found in further research.


Material and methods

The shells were collected by sieving the cave river deposits by 1x1 and 2x2 mm width of the mesh sieves. The material from the smaller meshed sieve was then brought to the laboratory and dried. After that it was again put into water and the floating shells were collected by a strainer and small brush. The measurements were carried out by means of CETI stereo microscope and an eye-piece micrometer, and photographs were made with camera system with a digital adapter. The material is stored in the National Museum of Natural History, Sofia, Bulgaria.

Abbreviations used: H - shell height, W - shell width, AH - aperture height, AW - aperture width, LH - last whorl height.

Table 1. Shell measurements of *Belgrandiella hubenovi* n. sp. (for abbreviations see “Materials and Methods”)

N	H	W	AH	AW	LH	W/H	AH/H	AW/AH	LH/H
1	1.49	0.76	0.54	0.54	1.06	0.51	0.37	1.00	0.71
2	1.49	0.89	0.56	0.56	1.02	0.60	0.38	1.00	0.69
3	1.67	0.89	0.63	0.59	1.12	0.53	0.38	0.95	0.67
4	1.73	0.99	0.71	0.63	1.19	0.57	0.41	0.88	0.69
5	1.55	0.89	0.56	0.59	1.12	0.57	0.36	1.06	0.72
6	1.62	0.86	0.63	0.59	1.12	0.53	0.39	0.95	0.69
7	1.49	0.87	0.56	0.59	1.09	0.59	0.38	1.06	0.73
8	1.68	0.87	0.63	0.59	1.19	0.52	0.37	0.95	0.71
Average	1.59	0.88	0.60	0.59	1.11	0.55	0.38	0.98	0.70

Fig. 1. Front and side view of the holotype of *Belgrandiella hubenovi* n. sp.

Results and discussion

Genus *Belgrandiella* Wagner, 1927

Type species: *Belgrandia kusceri* Wagner, 1914

Belgrandiella hubenovi n. sp.

Material examined: 9 shells, 09.03.2013, the Water cave (Vodnata) near village of Musina, north Bulgaria, N43° 13' 14.9" E25° 25' 39.6", 198 m alt., D. Georgiev leg.

Holotype: H = 1.49 mm, W = 0.76 mm, AH = 0.54 mm, AW = 0.54 mm, LH = 1.06

Paratypes: 3 shells, National Museum of Natural History, Sofia, Bulgaria, 5 shells coll. D. Georgiev.

Locus typicus: Water cave near village of Musina, north Bulgaria, N43° 13' 14.9" E25° 25' 39.6", 198 m alt.

Differential diagnosis: By its bent periostome the new species differs from all known Bulgarian species from the genus *Belgrandiella* with an exception of *B. hessei* – another stygobiont known from Temnata Dupka cave near Lakatnik town,

about 167 km to the west. From this species *B. hubenovi* n. sp. differs by its smaller size (H max = 1.73 mm versus 2.3 mm in *B. hessei*), and not so thick and developed aperture lip.

Description: Shell: The shell is ovate-conic to ovate-cylindrical, white, whorls are 3-4, regularly growing with, shining surface, with deep suture, fine growth lines sometimes forming small, irregular smooth ribs. The umbilicus is slit-like, the aperture is egg-shaped with sharp periostome which is bent to the outside from the lateral view. The operculum and the soft body are not known. H = 1.49-1.73 mm, W = 0.76-0.99 mm, AH = 0.54-0.71 mm, AW = 0.54-0.63 mm, LH = 1.02-1.19 mm, W/H = 0.51-0.60, AH/H = 0.36-0.41, AW/AH = 0.88-1.06, LH/H = 0.67-0.73.

Etymology: Named after Assoc. Prof. Dr Zdravko Hubenov (National Museum of Natural History, Sofia, Bulgaria) who contributed so much in the studies of the Bulgarian aquatic malacofauna and paid me attention of some springs as potential localities of freshwater snails.

Distribution: Known only from the type locality, possibly local endemic species.

Habitat and ecology: A stygobiotic species found only as empty shells in the river deposits at the entrance of Vodnata cave near village of Musina. This cave can be explored only by boat and experienced divers for searching living snails on the bottom substrate of the inner stream.

The systematic position of the genus *Belgrandiella* Wagner, 1927 is unclear. It is referred by various authors to different family groups as Hydrobiidae, Orientalinidae or Belgrandiellidae (Radoman, 1983; Kabat & Hersler, 1993; Angelov, 2000; Arconada & Ramos, 2003). The genus consists of minute Risoid snails with different shell shapes, mainly ovoid-conical but also conical or ovoid and different penis morphology (simple long,

simple short, conical or triangular, or same shape but with one small lobe on the left side). It can be supposed (as for other extremely subterranean genera, for example *Bythiospeum* Bourguignat, 1882) that *Belgrandiella* is a complex of more still undefined genera. However it is a very hard task and sometimes is impossible for the scientists to reach the habitats of some of the stygobiotic species.

References

- Angelov, A. 1959. Neue Gastropoden aus den unterirdischen Gewässern Bulgariens. Archiv für Molluskenkunde, 88(1/3): 51-54.
- Angelov, A. 1972. Neue Hydrobiidae aus Höhlengewässern Bulgariens. Archiv für Molluskenkunde, 102(1/3): 107-112.
- Angelov, A. 1976. Ein neuer Vertreter der Gattung *Belgrandiella* A. Wagner, 1927 (Gastropoda, Hydrobiidae) von Grundwassern Bulgariens. Acta Zoologica bulgarica, 4: 78-80.
- Angelov, A. 2000. Mollusca (Gastropoda et Bivalvia) aquae dulcis, catalogus Faunae Bulgaicae. – Pensoft & Backhuys Publ., Sofia, Leiden, 54 p.
- Arconada, B., Ramos, M. 2003. The Ibero-Balearic region: one of the areas of highest Hydrobiidae (Gastropoda, Prosobranchia, Rissooidea) diversity in Europe. Graellsia, 59 (2-3): 91-104.
- Georgiev, D. 2011a. Check list of the Bulgarian minor freshwater snails (Gastropoda: Rissooidea) with some ecological and zoogeographical notes. Zoonotes, 24: 1-4.
- Georgiev, D. 2011b. A New Species of *Belgrandiella* (Wagner 1927) (Mollusca: Gastropoda) from Caves in Northern Bulgaria. Acta Zoologica bulgarica, 63 (1): 7-10.
- Georgiev, D. 2011c. New species of snails (Mollusca: Gastropoda: Rissooidea) from cave waters of Bulgaria. Buletin Shkenkor, Ser. Shkenkat Natyore, 61: 83-96.
- Georgiev, D. 2011d. New localities of four Bulgarian Hydrobiidae species (Gastropoda: Rissooidea). Zoonotes, 24: 1-4.
- Georgiev, D. 2013. Catalogue of the stygobiotic and troglophilous freshwater snails (Gastropoda: Rissooidea: Hydrobiidae) of Bulgaria with descriptions of five new species. Ruthenica, 23 (1): 59-67.
- Glöer, P., Georgiev, D. 2009. New Rissooidea from Bulgaria (Gastropoda: Rissooidea). – Mollusca, 27 (2): 123-136.
- Hubenov, Z. 2005. Malacofaunistic diversity of Bulgaria. In: Current state of Bulgarian biodiversity – problems and perspectives, Petrova A. (ed.) Bulgarian Bioplatform, Sofia, 199-246. (in Bulgarian).
- Hubenov, Z. 2007. Fauna and zoogeography of marine, freshwater, and terrestrial mollusks (Mollusca) in Bulgaria. In: Fet, V., A. Popov (eds.). Biogeography and ecology of Bulgaria. Springer, Dodrecht, 141-198.
- Kabat, A., Hersler, R. 1993. The Prosobranch family Hydrobiidae (Gastropoda: Rissooidea): Review of classification and supraspecific taxa. Smithsonian Contributions to Zoology, 547: 1-94.
- Pintér, L. 1968. Eine neue Wasserschnecke aus Bulgarien. Archiv für Molluskenkunde, 98 (1/2): 61-63.
- Radoman, P. 1983. Hydrobioidea a superfamily of Prosobranchia (Gastropoda). I. Systematics. Monographs 547, Serbian Academy of Sciences and Arts, (Department of Science), 256 pp.
- Wagner, A. 1927. Studien zur Molluskenfauna der Balkanhalbinsel mit besonderer Berücksichtigung Bulgariens und Thraziens, nebst monographischer Bearbeitung einzelner Gruppen. Annales Zoologici Musei Polonici Hist. Nat., 6 (4): 263-399.